

NEW ROUTES TO HIGHLY EFFICIENT COPPER(I)-BASED DYE-SENSITIZED SOLAR CELLS: N^NX LIGANDS AS ANCILLARY LIGANDS

Frederik J. Malzner

(Department of Chemistry, University of Basel, Spitalstrasse 51, Basel, Switzerland, frederik.malzner@unibas.ch)

Renewable energy sources are the answer to save the world from drastic climate changes. The most promising one is photovoltaics. Today, photovoltaic modules are based on silicon but also consist rare elements. A very good alternative are dye-sensitized solar cells, which had its breakthrough in 1991 [1]. The most efficient cells contain ruthenium which is also rare on earth.

Our group focuses its research on the application of copper(I) complexes in dye-sensitized solar cells [2]. A dye-sensitized solar cells consists of different components, which can be optimized to yield higher efficiencies.




Figure 1: Structure of a dye-sensitized solar cell with zoom-in example for investigated copper(I) dye-sensitized solar cells.

The most important part of a dye-sensitized solar cell is the dye. It consists of two parts – the anchoring and the ancillary ligand. A series of heterocyclic N^NX ancillary ligands has been examined for the first time in copper(I) dye-sensitized solar cells. The relatively high efficiencies show the huge potential of copper(I) dye-sensitized solar cells [3].

[1] B. O'Regan, M. Grätzel, *Nature*, **1991**, 353, 737.

[2] C. E. Housecroft, E. C. Constable, *Chem. Soc. Rev.*, **2015**, 44, 8386.

[3] F.J. Malzner, M. Willgert, C. E. Housecroft, E. C. Constable, *manuscript in preparation*.